

TRANSYLVANIAN

REVIEW

SPECIAL ISSUE

VOL XXIV, No. 10, SPECIAL ISSUE, 2016



Transylvanian Review

Centrul de Studii Transilvane | str. Mihail Kogalniceanu nr. 12-14, et. 5, Cluj-Napoca

Email: transylvanianreview@gmail.com / WebSite: <http://www.centruldestudiitransilvane.ro/>

Online Submission System: <http://transylvanianreview.org/>

The Effects of Bio-Slurry Fertilizer on *Sorghum bicolor* and *Centrosema pubescens* Planted with Inter-Cropping

^{*1}Budiman Nohong, ²Tanrigiling Rasyidand ³Muhammad Yusuf

^{*1}Department of Forage Crops and Grassland Management, Faculty of Animal Husbandry, Hasanuddin University, Makassar, Indonesia 90245.

²Department of Social Economic, Faculty of Animal Husbandry, Hasanuddin University, Makassar, 90245, Indonesia

³Forestry extension agencies Bone regency South Sulawesi Indonesia.

Abstract

The aim of this study was to determine the effect of Fertilizer bio-slurry (urine-cow) against the production of dry matter, crude protein and competitiveness of plants Sorghum (*Sorghum bicolor*) and Centro (*Centrosema pubescens*) planted with monocultures (single) and poly (intercropping). The experiment was arranged in a factorial design consisting of three cropping systems, two levels of fertilizers and three replications. Cropping systems consist of sorghum and centro each planted single (sole) and sorghum-centro intercropping (2:2). Plants were fertilized by 0 liter of bio-slurry/ha and 25,000 liters of bio-slurry/ha (217.5 kg N/ha, 197.5 kg P₂O₅/ha and 230 kg K₂O / ha), or equivalent to 125 ml/ polybag (1, 1 g N/polybag, 1.0 g P₂O₅ /polybag and 1.2 g K₂O). Polybags in a size of 35 x 35 cm were filled by clay loam with ultisol texture weighing 10 kg on the field capacity condition. As many as six polybags were planted with eight sorghum grains; six polybags were planted with eight seeds of centro; while the other six polybags were planted with four sorghum grains and four centroseeds. The results showed that the use of bio-slurry markedly increase the production of dry matter and crude protein content of Sorghum and Centro. Sorghum fertilized by bio-slurry is more dominant indicated by its higher relative crowding coefficient (RCC), competitive ratio (CR) and positive sign on aggressiveness. Sorghum is grown with a more aggressive manner in utilizing resources than centro because it has greater competitiveness than centro.

Keywords: bio-slurry, centro, competitiveness, intercropping, sorghum.

^{*}Corresponding author: Department of Forage Crops and Grassland Management, Faculty of Animal Husbandry, Hasanuddin University, Makassar, Indonesia 90245.

Introduction

Intercropping is the cultivation of two or more crops or crop species, or genotypes, growing together at the same time in the same field (Ouma and Jeruto, 2010 and Brooker *et al.*, 2015). Sorghum has a good quality if it is grown along with legumes that propagates (Patel and Rajagopal 2003). Cereal-legume intercropping appears to be a useful component of ecological intensification approach to produce more food per unit resource to achieve positive social outcomes without negative effects on the environment (Dore *et al.*, 2011 and Hochman *et al.*, 2011). Intercropping of cereal and legumes contributes to maintain and improve soil fertility (Tsubo *et al.*, 2005). Legumes can transfer N for serealia during the growing period, and N is an important resource for serealia (Shen and Chu, 2004). Planting legumes and serealia altogether for food production is not only popular among farmers in the tropics, who produce most of the food in developing countries, but also it is developing into warm regions in the subtropics (Fujita *et al.*, 1992).

The main obstacles in improving and maintaining productivity and quality of forage on marginal land include lack of availability of nitrogen and water to support plant life. Meanwhile, the superior agricultural system has a superior specification of nitrogen (N), a demand that cannot be provided by reserved N substance in the soil. Consequently, an additional input is required (Senbayram *et al.*, 2016). However, N application will continue to accelerate the decrease of the non-renewable energy resources used in the production of fertilizers (Bohloul *et al.*, 1992), and the applications of organic and inorganic nitrogen fertilizers lead to a risk of inevitable ecological damages and a decline in production which make the cost of agricultural production tend to increase due to the decrease in soil pH of 0.5 units, in main crop production areas for two decades, and they have been attributed to excessive applications of N fertilizer (Guo *et al.*, 2010).

The agricultural system relying heavily on land reservation to fulfil the needs of N plants is unable to stay effective on producing high yields of crops (Stevenson, 1982). Except for legumes, they have the ability to fix their own N. Therefore, symbiotic N₂ fixation of legumes is widely utilized to increase the supply of N from the system. In addition to meeting the needs of N, legumes are able to facilitate the N intake from the surrounding plant species (Pirhofer-Walzl *et al.*, 2012).

Nowadays, a large number of unfortunate farmers cannot afford to buy fertilizers in sufficient quantities due to their high prices (Sisworo, 2006). Synthetic fertilizers are categorized as expensive commodities whose price is predicted to increase every year due to the decrease of their availability and the rising cost of fuel (Gensch *et al.*, 2011). This increase is assumed to bring serious implications for serealia grain production, particularly in industrialized countries located in subtropical areas as the

main serealia producers (Preston, 2000). Due to the rising costs of chemical fertilizers and their inefficient role at producing long-term sustainable production, then, the application of organic fertilizers such as bio-slurry (cow urine) to increase the maximum productivity in a sustainable manner is required and to provide better quality of the soil (Raj *et al.*, 2014).

Bio-slurry is an organic source obtained as a product after the generation of biogas from the fermentor. Bio-slurry removed from the reactor retains all the basic nutrients available in food sources, thus making bio-slurry as a potential organic fertilizer (SNV 2012), an environmentally friendly, inexpensive and renewable nutrient sources (Islam *et al.*, 2010), as well as a nutrient content resource in an available form (Bonten *et al.*, 2014). Nutrients in urine are available with a similar formulation to that of ammonia and urea resulting in comparable results for plant growth (Gensch *et al.*, 2011).

Efforts should be taken to build sustainable agricultural systems i.e. by optimizing the use of all external inputs that come from outside the system (Sanchez, 1994). One of the ways to perform the efforts is by supplying relatively inexpensive nutrients to utilize Biological Nitrogen Fixation (BNF). It is inarguable that BNF utilization is one of the N sources which is necessary in sustainable agricultural systems (Danso, 1995). In addition to BNF system, utilization of bio-slurry as an N source is also important in agricultural systems. BNF and bio-slurry utilization as source of nutrients is a truly renewable agricultural system as a whole. Such practices would pave the way to reduce the use of external inputs and improve sustainability among organic farmers in developing countries (Devakumar *et al.*, 2014). This study aims to determine the effect of liquid bio-slurry Fertilizer (cattle-urine) against the production of dry matter, crude protein and competitiveness of Sorghum (*Sorghum bicolor*) and Centro plants (*Centrosema pubescens*) planted in monoculture (single) and polyculture (inter-cropping).

Materials and Methods

This study was conducted at the Faculty of Animal Husbandry Research Station, University of Hasanuddin, Makassar, Indonesia (5°10' S, 119°20' E) from January 2016 to March 2016. During this study, the averages of daily temperature, humidity and rainfall were 27°C, 85 %, and 390 mm, respectively. Clay loam with ultisol texture was utilized.

The experiment was arranged in a factorial design consisting of three cropping systems, two levels of fertilizers and three replications. Cropping systems consist of sorghum and centro each planted single (sole) and sorghum-centro intercropping (2:2). Plants were fertilized by 0 liter of bio-slurry/ha and 25,000 liters of bio-slurry/ha (217.5 kg N/ha, 197.5 kg P₂O₅/ha and 230 kg K₂O/ha), or equivalent to 125 ml/ polybag (1, 1 g N/polybag,

1.0 g P₂O₅/polybag and 1.2 g K₂O/polybag). Polybags in a size of 35 x 35 cm were filled by clay loam with ultisol texture weighing 10 kg on the field capacity condition. As many as six polybags were planted with eight sorghum grains; six polybags were planted with eight seeds of centro; while the other six polybags were planted with four sorghum grains and four centro seeds. Each seed was individually planted at a depth of 3 cm. To get the required plant population, thinning the plants is conducted by uprooting some plants that are not needed and leaving four plants/polybag for sorghum and centro treatment which were planted single (monoculture); and for sorghum intercropping with centro, each was left with two sorghum/polybag and two centro/polybag. Applying fertilizer is done by spraying it around the plants. Watering and weeding were also conducted if necessary.

Data Collection

At harvest time, all the plants were harvested 5 cm above the ground by using scissors and weighed to determine the fresh weight. Thus, fresh samples chopped and dried in an oven at a temperature of 55°C for 3 days, and later they were weighed to determine the dried ingredients. Dried matter production = dry matter content x fresh weight production. All samples were milled (1 mm) with a grinding machine. This sample was used to determine the total nitrogen (N) by Kjeldahl method (AOAC, 2005), and crude protein (CP) was calculated by N x 6.25.

The competitiveness index was calculated using the competition index. Competition index has been widely used to determine the level of competition between species planted with intercropping system.

Competition index used in this study was relative yield Index (RY), relative yield total (ryt), relative crowding coefficient (RCC) and the aggressivity index (AI).

Relative yield (RY) and relative yield total (RYT) were counted by using Wiley's formulation (1979) as follows:

$$RY_{\text{Sorghum}} = \text{DMYss}/\text{DMYsi},$$

$$RY_{\text{Centrossema}} = \text{DMYcs}/\text{DMYci}$$

$$\text{RYT} = RY_{\text{Sorghum}} + RY_{\text{Centro}}$$

Where: dry matter production of sole planted Sorghum plant.

DMYss = dry matter production of sorghum planted in

monoculture.

DMYcs = dry matter production of centro planted in monoculture.

DMYsi = dry matter production of sorghum planted by

means of intercropping with centrol.

DMYci = dry matter production of centro grown by means of

intercropping with sorghum.

If RYT value is greater than one, it indicates that the species requires different resources to avoid competition.

Meanwhile, if RYT value is less than one, it indicates a mutually antagonistic, and if RYT value equals to one, it indicates that the component species share the same limited resources (Harper, 1977).

LER was utilized as a criterion for the benefits of mixed stands. In particular, LER showed intercropping efficiency in using resources from the environment compared to single cropping (Mead and Willey 1980). If LER is greater than one, intercropping tends to be profitable and productive. On contrary, if LER is less than one (<1), intercropping affects negatively on the growth and crops yield planted in a mixture (Ofori and Stern 1987). This is a complementary indicator in which LER was calculated in reference to techniques proposed by Wiley and Osiru (1972) as follows:

$$\text{LER} = (\text{YSC}/\text{YSS}) + (\text{YCS}/\text{CC}),$$

Where:

YSC = Result of sorghum crops when intercropped (poly) with

plants centro.

YCS = Result of centro crops when intercropped (poly) with

sorghum.

YSS = Result of single grown sorghum (monoculture).

YCC = Result of single grown sentro (monoculture).

The relative crowding coefficient = RCC or K is a measure of the relative dominance of one species compared to other species in mixture (Willey and Rao 1980), calculated by the formulation of: RCC or K = (K Sorghum x K Centro), where K sorghum = YSI x Zcp / [(Yss - YSI) x ZSP], and K Centro = YCI x ZSP / [(YCS - Yci) x Zcp], where ZSP (= 2) and Zcp (= 2) are the proportion of sorghum and centro in mixture. Plant components that have a higher coefficient is considered to be more dominant. If the coefficient value of certain plants is less than one, the crop yields become less; if the coefficient value is equal to one, the result is the same, whereas when the coefficient value is greater than one, the result is more than it is "expected" (Willey and Rao 1980).

Competition ratio = CR was measured to determine the competition level between plant crops in the cropping system (Bhatti 2006). This ratio reflects the individual competition of the two components of LER plants while taking into an account the proportion of crop planted.

Competition ratio of sorghum and cowpea in the mixture was calculated by the formula proposed by Willey *et al.*, (1980) as follows: CR = sorghum (sorghum LER / LER centro) (Zc / Zs), CR centro = (LER centro / LER sorghum) (Zs / Zc). Zs (= 2) and Zc (=2) are the proportion of sorghum and centro in the mixture.

Aggressivity indices = AI are a measure of the relation of competition between two crops in a mixed cropping (mixtures) (Willey, 1979). This is discussed by Dhima *et al.*, (2007) as follows: AI Sorghum = (YSI / Yss x Zp) - (YCI / YCS x Zc) and AI Centro = (YCI / YCS x Zc) -

(YSI / Yss x Zs). So, if AI sorghum is 0, both of these plants are equally competitive, if AI sorghum is positive, then the AI sorghum is more dominant; while if it is negative, then the sorghum is weaker.

Statistical Analysis

The effects of treatment on dry matter production and crude protein content were calculated by using SPSS software (SPSS Ver. 16.0, SPSS Inc. Chicago, Illinois). Then, LSD test was carried out to examine the different treatments. The probability level of $P \leq 0.05$ is considered to be statistically significant.

The competitiveness index was calculated by using the competition index. Relative yield = RY and relative yield total = RYT were calculated by using the formula proposed by Willey (1979). LER was calculated in accordance with (Willey and Osiru, 1972), relative crowding coefficient = RCC (Willey and Rao, 1980),

competitive ratio = CR is calculated by the formula Willey *et al.*, (1980) and aggressivity index = AI) is a ameasurement form of the competition relation between two plants in the mixed planting (polyculture) (Wiley, 1979)

Results

Effect of Cropping System on Dry Matter Yield and Crude Protein Content

The summary of statistical analysis of data for dry matter yield, crude protein content is shown in Table 1. Cropping system caused significant response ($P < 0.05$) on dry matter yield and crude protein content (Table 1). Dry matter yield of sole sorghum was significantly higher ($P < 0.05$) than the sorghum-centro intercropping and sole centro. Crude protein content of sole centro was significantly higher ($P < 0.05$) than the sorghum-centro intercropping and sole sorghum.

Table 1: Effects of cropping systems on dry matter yield and crude protein content.

Parameters	Cropping system	Fertilizer		Average	Probability	SEM
		P0	P1			
Yield (g/pot)	Sole crop of Sorghum	17.36	24.77	21.07 ^a	0.000	1.099
	Sole crop of Centro	6.66	13.33	10.00 ^c	0.000	1.099
	Sorghum-centro (2:2)	12.82	18.80	15.81 ^b	0.000	0.457
Crude protein (%)	Sole crop of Sorghum	6.11	7.75	7.26 ^c	0.000	0.484
	Sole crop of Centro	11.77	12.70	11.79 ^a	0.000	0.000
	Sorghum-centro (2:2)	6.86	7.84	8.69 ^b	0.000	0.457

^{abc}: the different superscripts within the column indicate differences ($p < 0.05$).

Effect of Bio-Slurry Fertilizer on Dry Matter Yield and Crude Protein Content

The effect of bio-slurry fertilizer on the dry matter yield crude protein content of sorghum-centro intercropping system is presented in Table 2. Application of bio-slurry fertilizer increased ($P < 0.05$) dry matter yield

and crude protein content of sorghum sole and centro sole as well as sorghum-centro intercropping, compared without fertilizer (control). The increased dry matter and crude protein content as a result of bio-slurry fertilizer were 54% and 14%, respectively.

Table 2: Effects of bio-slurry Fertilizer on the production of dry matters and crude protein Content.

Parameters	Fertilizer	SS	CS	S:C (2:2)	Avrg	Prob.	SEM
Yield (g/pot)	P0	17.36	6.66	12.82	12.28 ^b	0.000	0.969
	P1	24.77	13.33	18.80	18.97 ^a	0.000	0.859
Crude protein (%)	P0	6.11	11.77	6.86	8.25 ^b	0.048	0.403
	P1	7.75	12.70	7.84	9.43 ^a	0.048	0.357

SS = Sole crop of Sorghum, CS = Sole crop of Centro, S:C = sorghum-centro intercropping (2:2),

SEM = Standard Error of Mean.

Competitiveness

Relative Yield (RY) and Relative Yield Total (RYT)

Relative yields (RY) and (relative yield total (RYT) of centro and sorghum in intercropping systems are

presented in Table 3. In Table 3, it appears that RY sorghum (RYS) and RY centro (RYC) at P0 and P1 value are less than one (< 1). The RYT values as observed were 0.99 (P0) and 0.89 (P1) in treatments (2:2).

Tabel 3: Pengaruh pemberian pupuk bio-slurry terhadap hasil relatif sorgum (RYS), hasil relatif centro (RYC) dan hasil relatif total (RYT).

Fertilizer	YSS	YSI	YCS	YCI	RYS	RYC	RYT
P0	17.36	11.36	6.66	2.29	0.65	0.34	0.99

P1	24.77	17.49	13.33	2.48	0.71	0.19	0.89
----	-------	-------	-------	------	------	------	------

YSS = yield sorghum as sole crops, YSI = yield sorghum as intercrop, YCS = yield Centro as sole crops, YCI = yield Centro as sole intercrops.

Land Equivalent Ratio (LER)

The effects of bio-slurry Fertilizer on the value of land equivalent ratio (LER) in intercropping system of sorghum-

centro (2: 2) are presented in Table 4. The value of land equivalent ratio (LER) is smaller than one (<1).

Tabel 4: Pengaruh pemberian pupuk bio-slurry terhadap Land equivalent ratio (LER), Relative Crowding Coefficient (RCC), Competitive ratio (CR) dan Indeks Agresivitas (AI).

Fertilizers	LER		LER TOTAL	RCC		CR		AI	
	S	C		S	C	S	C	S	C
P0	0.65	0.34	0.99	4.61	0.64	1.91	0.52	0.15	-0.15
P1	0.71	0.19	0.90	4.24	0.69	3.74	0.27	0.27	-0.27

S = Sorghum, C = Centro.

Relative Crowding Coefficient (RCC)

The impact on competition and the advantages of intercropping are determined by calculating the coefficient of relative crowding. The values of relative crowding coefficient (RCC) of sorghum and centroplanted with intercropping system are presented in Table 4. The RCC value of sorghum is higher than that of centro of both treatments i.e without Fertilizer (P0) and with Fertilizer (P1).

Competitive Ratio (CR)

The effects of bio-slurry fertilizer on the competitive ratio between sorghum and centro are presented in Table 4. The competitive value in the fertilizer application of 3.74 (P1) indicates that this competitive ratio of sorghum-centro intercropping is the highest and the value of 1.91 indicates the lowest competitive ratio in the treatment without fertilizers (P0).

Aggressivity Index (AI)

The experiment showed that aggressivity index at intercropping sorghum-centro (2:2) showed that the two species have the same numerical value, but different signs. Sorghum aggressiveness index (+0.15) and centro (-0.15) in the treatment without fertilizer (control). Application of bio-slurry fertilizer increased indices aggressiveness of sorghum (+0.27) and centro of (-0.27).

Discussion

Differences in forage yield between of sorghum and centro may be attributed by different in species. Sorghum growing faster and capable of producing a higher dry matter rather than centro. The overall results indicated that dry matter yield of sorghum grown in association with centro was lower than the sole crop of sorghum which was probably the result of plant competition for the nutrients. According to Buxton and Fales (1993) that the interplant competition usually includes competition for soil water,

available nutrients, and solar radiation. Siddig *et al.* (2013) reported that dry matter weight of sorghum was influenced due to sorghum-ground nut intercropping. In contrast, other researchers have reported that dry matter yield of sorghum greater when grown with groundnuts (Krishna *et al.*, 1998 and Ayisi *et al.*, 2004). Application of bio-slurry increased response ($P < 0.05$) on dry matter yield of sorghum sole, centro sole and sorghum-centro intercropping. The responses are due to the addition of soil nutrients derived from bio-manure slurry. According Mburu (2012) that the urine (slurry) is a source of macro nutrients such as NPK and other elements needed by plants. Increasing the level of slurry nitrogen presumably increased the availability of soil nitrogen, and that of other macro- and micronutrients, which might have enhanced meristematic growth and resulted in higher fodder yield (Islam *et al.*, 2010). Effect of bio-slurry on the growth and production of crops has been reported by several investigators. The utilization of cow-bio-urine promotes plant growth and yield in crops of eggplant (Cardoso *et al.*, 2009), corn (Devakumar *et al.*, 2014), carrots and okra (Muhmood *et al.*, 2015) and the elephant grass and Setaria (Nuriyasa *et al.*, 2012).

Application of bio-slurry increased response ($P < 0.05$) on dry matter yield of sorghum sole, centro sole and sorghum-centro intercropping. The responses are due to the addition of soil nutrients derived from bio-manure slurry. Urine (slurry) is a source of macro nutrients such as NPK. Increasing the level of slurry nitrogen presumably increased the availability of soil nitrogen, which might have enhanced meristematic growth and resulted in higher fodder yield (Islam *et al.*, 2010 and Mburu, 2012). Bio-slurry can play an important role as a source of nutrients for crop production. Compared to manure, nutrients in bio-slurry (especially nitrogen) are more readily available, which means that bio-slurry can have a larger fertilization effect in short term (Bonten *et al.*, 2014). Effect of bio-slurry on the growth and production of crops has been reported by several investigators.

The utilization of cow-bio-urine promotes plant growth and yield in crops of napier grass and *Setaria* (Nuriyasa *et al.*, 2012), eggplant (Cardoso *et al.*, 2009), corn (Devakumar *et al.*, 2014), carrots and okra (Muhmood *et al.*, 2015), legume crop pueri (Mudhita *et al.*, 2016). Application of bio-slurry increased response ($P < 0.05$) on dry crude protein content of sorghum sole, centro sole and sorghum-centro intercropping. The applications of dry bioslurry o increase crude protein content in maize (Rahman *et al.*, 2008) and grass (Mikled *et al.*, 2002). Sorghum-centro intercropping improves the overall quality of forage quality, because legumes play an important role in the supply of nitrogen. Legumes have the ability to fix N_2 from the air so that they have a higher protein content. As with other legums, centro can improve soil fertility through nitrogen fixation, and nitrogen fixation by legumes begins with the formation of nodules. The results of this study indicate that the centro is able to form nodules. It is able to fixate the equivalence of 67 kg N/ha and can transfer N at 5 kg/ha on the mixed crops of guinea-centro (Reynolds, 1982). Planting grass along with legumes is helpful in producing grass since Fertilizer is not necessary (Castillo *et al.*, 2003). Crude protein content of the non-legume crops can be increased if planted with plants from the legume family (Eskandari *et al.*, 2009).

According to De Wit (1960), that if the values of RY (species A) and RY (species B) are both less than one (< 1), it is evident that there is a mutual antagonism between the competitors. If one species is reduced, then there are many resources available for other species. This shows that there is a mutual antagonism between the centro and sorghum in intercropping. According to Williams and McCarty, (2004) if one species is reduced by 30% in the mixed crops, then the RY value is 0.7; however, If the species has experienced a 30% increase in the mixed crops, then the RY value is 1.3. Relative yield total = RYT (Table 3) showed that the values of RYT on the treatments without fertilizer (P0) and with fertilizer (P1) were less than one (< 1).

According to Silvertown and Lovett Doust (1993), when the RYT value is 1.0, then there is a competition between species to get the same resources; however, if RYT value is greater than one (> 1), then there are plant species that avoid competition to occur; and if the RYT value is less than one (< 1), there is a mutual antagonism between plant species. This shows that sorghum and centro planted together in polybag have decreased their dry matter production. The decline is due to competition for limited resources that exist in the growing media.

The land equivalent ratios (LERs) were assessed in terms of dry matter production of two treatments used. Results showed that the LER values of dry matter production for both sorghum-centro mixtures substantially less that of their corresponding monoculture. Since the idea of land equivalent ratio in most cases, is the most important comparison between yield of the main crop in

mixture and its yield in pure stand, it is rather expected that the combination of component species in mixture will be more productive than the species grown as a sole crops. The result obtained were not in accordance with the definition of land equivalent ratio in that the combination of component species in the mixture were more productive than the same species when grown as sole crops. The LERs ratio, in P0 (0.99) and P1 (0.90) treatments, were less than 1, the sole which interpreted as disadvantage of mixture over sole crop. Because it is less than one (< 1), it will give a negative effect on growth and yield (Ofori and Stern 1987). This condition indicates that the cropping patterns of sole centro or sole sorghum will provide greater results than those by the intercropping of sorghum-centro. According to Mead and Willey (1980), LER shows the efficiency of intercropping in using resources from the environment is greater compared that of sole planting. Ghosh (2004) reported that the Veth-barley mixture and mix Veth-triticale showed losses compared with pure stands.

The aggressiveness of one species towards another demonstrates by the value of relative crowding coefficient (RCC). Plant components that have a higher relative crowding coefficient (RCC) value is more dominant. To determine if there are any profits from the yields of cropping systems, the coefficient values of both plant components must be obtained. If the value of relative crowding coefficient (RCC) of the two species is of the same value, then it means that the planting system does not have any advantages. Intreatments (P0) and (P1), sorghum appeared to be highly dominant as it had higher value of RCC than the intercrops in different intercropping sorghum system (Table 4).

If the value of the RCC is less than one, then the planting system has some weaknesses/shortcomings; however, and if the value of RCC is more than one (> 1), then the planting system has advantages or benefits. In this study, the biggest advantage recorded is from sorghum which was treated both by fertilizers (P1) and without Fertilizer (P0) their RCC values were 4.24 and 4.61 respectively. It can be inferred that the intercropped sorghum utilized the resources more competitively and centro appeared to be dominated.

The effects of bio-slurry fertilizer on the competitive ratio between sorghum and centro are presented in Table 4. The competitive value in the fertilizer application of 3.74 (P1) indicates that this competitive ratio of sorghum-centro intercropping is the highest and the value of 1.91 indicates the lowest competitive ratio in the treatment without fertilizers (P0). This indicates that the bio-slurry fertilizer in intercropping sorghum-centro (2:2) promotes the growth of sorghum and suppress the growth of centro. In all mixtures, a positive sign of sorghum showed that sorghum was dominant species in the mixture (Table 4). Intercropping sorghum-centro has CR 1.91 for sorghum and 0.52 for the centro. Application of bio-manure slurry

increases to CR sorghum 3.74 and decreased to CR centro 0.27. High competitive value ratio (CR) indicates high level of competition between plants intercropped. Ariel *et al.*, (2013) state that the more dense the composition of plants in the intercropping, the higher the competition

Aggressivity index is used to determine the magnitude of the competition relationship in mixed cropping systems (Willey, 1979). The aggressivity index (AI) of sorghum-centro intercropping fertilized by bio-slurry fertilizer (P4) and without fertilizer (P0) showed that sorghum has a positive sign (+) while centro has a negative sign (-) (Table 4). The positive sign on sorghum showed that it was dominant while the negative sign (-) on centro indicated that centro was being dominated (weak). Dhima *et al.*, (2007) state that the zero value of aggressivity index indicated that the plant components were equally competitive. If a plant has the same numerical value, the positive sign (+) indicates the dominant species (strong), while the negative sign (-) showed that a species is being dominated (weak). This shows that sorghum is more dominant than centro planted in polybags.

Conclusion

Fertilizer using bio-slurry increases the production of dry matter and crude protein content of sorghum and centro planted with a monoculture system and sorghum-centro planted with a polyculture system. Sorghum treated by bio-slurry fertilizer is more dominant as shown by its high values of the relative crowding coefficient, competitive rates and positive sign of aggressivity. Sorghum has grown more aggressively than centro in utilizing resources.

References

- AOAC (2005). Official Methods of Analysis Association (18th. Ed.), Official Analytical Chemist. Association of Official Analytical Chemist, Washington, DC.
- Ariel, C.E., Eduardo, O. A., Benito, G.E., Lidia, G.(2013). Effects of two plant arrangements in corn (*Zea mays* L.) and soybean (*Glycine max* L. Merrill) intercropping on soil nitrogen and phosphorus status and growth of component crops at an Argentinean Argiudol. American Journal of Agriculture and Forestry. Vol. 1(2):22-31.
<http://www.sciencepublishinggroup.com/j/ajaf> doi: 10.11648/j.ajaf.20130102.11
- Ayisi, K.K., M.S. Mpori and Van-den-Berg, J. (2001). Grain yield response and chilopartillus infestation in sorghum-cowpea intercrop management. *South African J. Pl. Soil.*, 18(4): 39-42.
- Bhatti, I.H., Ahmadi, R., Jabbar, A., Nazir, M.S., & Mahmood, T. 2006. Competitive Behaviour of Component Crops in Different Sesame-Legume Intercropping Systems. *Int. J. Agri. Biol.*, Vol. 8, No.2 :165–167. <http://www.fspublishers.org>
- Bohlool, B. B., J. J. Ladha, D P. Carrity and George, T (1992). Biological nitrogen fixation for sustainable agriculture: A perspective. *Plant and Soil* 141: 1–11.
<http://link.springer.com/article/10.1007%2FBF00041307>
- Bonten, L.T.C., K.B. Zwart, R. P. J. J Rietra, R. Postma and de Hass, M.J. G.(2014). Bio-slurry as fertilizer. Wageningen University.
<https://www.researchgate.net/publication/268148367>. DOI: 10.13140/2.1.2811.1040
- Brooker, W.W., E. Alison, Bennett, C. Wen-Feng, J. D. Tim, S.G. Timothy, D.H.Paul, H.Cathy, P. P. M. Iannetta, G. J.Hamlyn, J. K. Alison, L. Li, B. M. McKenzie, R. J. Pakeman, E. Paterson, S. Christian, J. Shen, G. Squire, C. A. Watson, C. Zhang, F. Zhang, J. Zhang and Philip J. W. (2015). Improving intercropping: a synthesis of research in agronomy, plant physiology and ecology. *New Phytologist*, 206: 107–117.
www.newphytologist.com. doi: 10.1111/nph.13132.
- Buxton, C.L. and Fales, S.L. (1993). Plant environment and quality. In: Fahey, Jr., G. C. (Ed.) *Forage Quality, Evaluation and Utilization*. ASA, CSSA and SSSA, Madison, WI.
- Cardoso, M.O., Olivera, A.P., Pereira, W.E. and Souza, A.P. (2009). Growth, nutrition and yield of eggplant as affected by doses of cattle manure and magnesium thermophosphate plus cow urine. *Horticultura Brasileira* 27: 307-313.
www.scielo.br/pdf/hb/v27n3/v27n3a08.pdf
- Castillo, E., T.E. Ruiz, R. Stuart, J. Galindo, J.L. Hernandez and Diaz, H.(2003). Effect of the protein-energetic supplementation on the performance of male bovines grazing natural pastures associated with a mixture of creeping legumes. *Cuban J. Agric. Sci.*, 37 (2): 143-147. www.feedipedia.org/node/1786
- Daniel, T.B., B. Lammert, Kropff, M J. (2004). Competition and crop performance in a leek–celery intercropping system. *Crop Sci* 41: 764–774.
- Danso, S.K.A. 1994. Sustainable Agriculture : the role of biological nitrogen fixing plants, *In: Nuclear Techniques in Soil-Plant Studies for Sustainable Agriculture and Environment Preservation*. Proceedings, International Symposium on Nuclear and Related Techniques in Soil-Plant Studies on Sustainable Agriculture and Environmental Presentation. Vienna, 17 -21 October 1994, Page 205–224).
https://inis.iaea.org/search/search.aspx?orig_q=RN:26056401
- de Wit, C.T. (1960). On Competition. *Agricultural Research Reports* 66(8) : 1 – 28. Centre for Agricultural Publishing and Documentation (PUDOC), Wageningen, The Netherlands.

- Devakumar, N., S. Shubha, G. G. E. Rao and Imrankhan. (2014). Studies on Soil fertility, Cow urine and Panchagavya levels on Growth and Yield of Maize. Proceedings, the 4th ISOFAR Scientific Conference. 'Building Organic Bridges', at the Organic World Congress 2014, 13-15 Oct., Istanbul, Turkey. Page 620 – 630
- Dhima, K. V., A.S. Lithourgidis, I. B. Vasilakoglou and Dordas, C. A. (2007). Competition indices of common vetch and cereal intercrops in two seeding ratio. *Field Crops Res.* 100: 249-256.
data2.xjlas.ac.cn:81/UploadFiles/sdz/.../intercropping/85.pdf
- Dore, T., K. Cadoret, D. Makowski, E. Malezieux, N. Munier-Jolain, M. Tchamitchian, Tittonell, P. (2011). Facing up to the paradigm of ecological intensification in agronomy: revisiting methods, concepts and knowledge. *Eur. J. Agron.* 34: 197-210. doi:10.1016/j.eja.2011.02.006.
- Eskandari, H. (2011). Intercropping of wheat (*Triticumaestivum*) and bean (*Viciafaba*): Effects of complementarity and competition of intercrop components in resource consumption on dry matter production and weed growth. *African Journal of Biotechnology* 10(77): 17755-17762.
<http://www.academicjournals.org/AJB>. DOI: 10.5897/AJB11.2250
- Fujita, K., G. Ofosu-Budu, and S. Ogata. 1992. "Biological nitrogen fixation in mixed legume-cereal cropping systems," *Plant and Soil*, Vol. 141, No. 1, pp. 155–175.
<https://www.researchgate.net/publication/225966474>. DOI: 10.1007/BF00011315
- Gensch, R., Miso, A., & Itchon, G. (2011). Urine as Liquid Fertilizer in Agricultural Production in the Philippines A Practical Field Guide. Xavier University Press. p.48 [Online] Available : Urine as liquid fertilizer (pdf)
- Giller, K.E. (2001). Nitrogen Fixation in Tropical Cropping Systems, (2nd ed)., CABI, Wallingford, p. 423
- Guo, J. Liu, X.J. Zhang, Y. Shen, J.L. Han, W. X. Zhang, W.F. (2010). Significant acidification in major Chinese croplands. *Science* 327 (5968), 1008–1010. doi: 10.1126/science.1182570.
- Harper, J.L. (1977). *Population Biology of Plants*. Academic Press, London.
- Hochman, Z., P.S. Carberry, M.J. Robertson, D.S. Gaydon, L.W. Bell, McIntosh P.C., (2011). Prospects for ecological intensification of Australian agriculture. *Eur. J. Agron.* 44: 109-123
- Islam, M.R., Rahman, S.M.E., Rahman, M.M., D.Hwan, Six, R.C. (2010). The effects of biogas slurry on the production and quality of maize fodder. *Turk J Agric For.* 34: 91–99. doi:10.3906/tar-0902-44.
- Krishna, A., S.V. Raikhelkar and Reddy, A.S. (1998). Effect of planting pattern and nitrogen on fodder maize (*Zea mays*) intercropped with cowpea (*Vignaunguiculata*). *Indian J. Agronomy*, 43(2): 237-240. [CAB Absts. 1998/08].
- Mburu, C. (2012). Urine as a fertilizer for lettuce grown in greenhouse condition. Degree programme in Environmental Engineering Tampereenammattikorkeakoulu Tampere University of Applied Sciences. Bachelor Thesis 42 pages.
<http://urn.fi/URN:NBN:fi:amk-2012102914699>
- Mead, R and Willey, R W (1980) *The concept of a 'land equivalent ratio' and advantages in yields from intercropping*. *Experimental Agriculture*, 16 (3). pp. 217-228. ISSN 0014-4797
- Mikled, C., J. Suchart and Potikanond, N. (2002). Utilization of fermented slurry as bio-fertilizer. Proceedings Biodigester Workshop March 2002.
<http://www.mekarn.org/procbiod/choke.htm>
- Muhmood, A., M. Abdul, N. Abid, S. Javid, S.S.H. Syah and Shah, A.H. (2015). Nutrients Uptake and the Yield of Okra and Carrot in Response to Bioslurry and Inorganic N Fertilizers. *International Journal of Plant & Soil Science*, 7(5): 297-305. www.sciencedomain.org
- Nuriyasa, I.M., N. N. Candraasih K., A. A. A. S. Trisnadewi, E. Puspani, W. Wirawan, (2012). Peningkatan Produksi Gajah (*Pennisetum purpureum*) dan Rumpun Setaria (*Setariasplendida* Stapf) Melalui Pemupukan Biourin. *Pastura* Vol. 2 No. 2 : 93 – 96.
<http://ojs.unud.ac.id/index.php/pastura/article/view/9029>
- Ofori, F. and Stern, W.R. 1987. Cereal-legume intercropping systems. *Advances in Agronomy*, 40:41-90.
- Ouma, G. and Jeruto, P. (2010). Sustainable horticultural crop production through intercropping: The case of fruits and vegetable crops: A review. *Agric. Biol. J. N. Am.*, 2010, 1(5): 1098-1105.
<http://www.scihub.org/ABJNA>. doi:10.5254/abjna.2010.1.5.1098.1105
- Patel, J. R. and S. Rajagopal. (2003). Nitrogen management of sorghum (*Sorghum bicolor*) and cowpea (*Vignaunguiculata*) forage under intercropping system. *Indian Journal of Agronomy*. 48 (1): 34 - 37.
- Pirhofer-Walz, K J. Rasmussen, H Høgh-Jensen, J Eriksen, K. Soegaard dan J. Rasmussen. 2012. Nitrogen transfer from forage legumes to nine neighbouring plants in a multi-species grassland. *Plant Soil*, 350: 71–84. DOI 10.1007/s11104-011-0882-z
- Preston, T. R. (2000). Livestock Production from Local Resources in an Integrated Farming System; a Sustainable Alternative for the Benefit of Small Scale Farmers and the Environment. Proceedings National Workshop-Seminar Sustainable Livestock Production on Local Feed Resources, Ho Chi Minh City, Vietnam January 18-20th.
<http://www.mekarn.org/sarpro/preston.htm>
- Rahman, S. M. E., Islam M. A., M.M. Islami, M. M. and Deog-Hwan Oh. Effect of Cattle Slurry on Growth, Biomass Yield and Chemical Composition of Maize

- Fodder. Asian-Aust. J. Anim. Sci. Vol. 21, No. 11 : 1592 – 1598. ajas.info/upload/pdf/21-221.pdf.
- Raj, A., K. J. Manoj and T. Pratap (2014). Cow Dung for Ecofriendly and Sustainable Productive Farming. International Journal of Scientific research, Vol. 3 No 10. 101 – 102. DOI : 10.15373/22778179
- Reynold, S. G. (1982). Contribution to Yield, Nitrogen Fixation and Transfer by Local and Exotic Legumes in Tropical Grass-Legume Mixtures in Western Samoa. Tropical Grass Legume in Samoa. Tropical Grasslands. Volume 16: 76–80.
- Sanchez , P.A. (1994). Tropical soil fertility research: towards the second paradigm. In : Proc. Of the 15 th Word Congress of Soil Science. International Society of Soil Science, Acapulco, Mexico Vol. 1. 66 – 88.
- Senbayram ,M., W. Christian, L. Annika, I. Johannes, S. Horst, K. Cengiz and Sarah, K.(2016). Legume-based mixed intercropping systems may lower agricultural born N₂O emissions. Energy, Sustainability and Society, 6:2, p:9.
- Shen, Q.R., Chu, G.X. (2004). Bi-directional nitrogen transfer in an intercropping system of peanut with rice cultivated in aerobic soil. Biol. Fertil. Soils. 40: 81-87.
- Siddig. A., Mohamed, Ali, Adam. A. Mohamed, Ali. H. Bahar and Abdulmohsin. R. Khairseed. 2013. Effects of Sorghum (*Sorghum bicolor* L.) Moench) and Groundnut (*Arachis hypogaea* L.) Intercropping on some soil chemical properties and crop yield under rain-fed conditions. *ARPJ Journal of Science and Technology*, 3:1.
- Silvertown, J. and Lovett Doust, J. (1993). Introduction to Plant Population Biology. Blackwell Scientific Publications, Oxford, UK.
- Sisworo, W. H.(2006). Swasembada Pangandan Pertanian Berkelanjutan Tantangan Abad Dua Satu : Pendekatan Ilmu Tanah Tanaman dan Pemanfaatan Iptek Nuklir. Badan Tenaga Nuklir Nasional.
- SNV, (2012). Promotion of Bio slurry for Increased Production in Organic Farming in Tanzania. www.ifoam.bio/.../banzi_bioslurry_and_organic_farming_snv....
- Stevenson, F.J. (1982). Origin and distribution of nitrogen in soil. In: Nitrogen in agricultural soils, F. J. Stevenson, Ed., 1–42. Madison, WI: ASA, CSSA, and SSSA.
- Tsubo, M. S. Walker and Ogindo, H. O. (2005). A Stimulation model of cereal-legume intercropping systems for semi-arid regions. Field Crops Research. 93: 10 - 22.
- Willey, R. W., Rao, M. R.(1980). A competitive ratio for quantifying competition between intercrops // Experimental Agriculture. – 1980, vol. 16, p. 117–125
- Willey, R.W. (1979). Intercropping – its importance and research needs. Part 1. Competition and yield advantages. Field Crop Abstr. 32,1–40.
- Willey, R.W. and Osiru, D.S.O. (1972). Studies on Mixtures of Maize and Beans (*Phaseolus-Vulgaris*) with Particular Reference to Plant Population. Journal of Agricultural Science 79: 517-529.
- Williams, A. C. and McCarthy, B.C. (2004). A new index of interspecific competition for replacement and additive design. Ecological Research. Vo. 16 : 29 – 40.